

PTO/PCT Rec'd 17 JAN 2001

AGROBACTERIUM-MEDIATED TRANSFORMATION  
OF TURFGRASS

This application claims priority to U. S. Provisional Application No. 60/093,163, filed July 17, 1998, the entirety of which is incorporated by reference herein.

5

**FIELD OF THE INVENTION**

This invention relates to the field of plant transformation methods. More specifically, an *Agrobacterium*-mediated method for transforming turfgrass is provided, as well as transgenic turfgrass produced by the method.

10

**BACKGROUND OF THE INVENTION**

Various scientific articles and patents are referred to in parentheses throughout the specification. These documents are incorporated by reference herein to describe the state of the art to which this invention pertains. Full citations of the scientific articles appear at the end of the specification.

15

Turfgrasses and turfgrass breeding are of significant economic importance worldwide. In recent years, traditional breeding programs have been augmented by molecular biological and recombinant techniques. However, similar to most monocotyledenous plants, turfgrasses have proven recalcitrant to tissue culture, transformation and regeneration procedures. Among these, *Agrobacterium*-mediated transformation of a turfgrass heretofore has not been accomplished.

20

25

Turfgrass transformation has been achieved using direct methods of DNA transfer, including protoplast transformation and particle gun bombardment.

30

- 2 -

Nonetheless, *Agrobacterium*-mediated transformation offers several advantages over particle gun bombardment or other means of direct gene transfer. These include stable transgene integration without rearrangement of either  
5 host or transgene DNA; preferential integration of the transgene into transcriptionally active regions of the genome; ability to transfer large pieces of DNA; and integration of low numbers of gene copies into plant nuclear DNA which is particularly important to minimize  
10 possible co-suppression of the transgene in later generations.

Until recently, *Agrobacterium*-mediated transformation was thought to be limited to dicotyledonous plants. However, Hiei et al. in 1994  
15 described efficient transformation of rice by *Agrobacterium*, and subsequently there have been convincing reports for maize, barley and wheat (Ishida et al., 1996; Tingay et al., 1997; Cheng et al., 1997; see also U.S. Patent No. 5,591,616 to Hiei et al). Numerous  
20 factors are of critical importance in *Agrobacterium*-mediated transformation of monocots, including the type and stage of tissue that is infected, the vector and bacterial strains used, plant genotype, tissue culture conditions, and the actual infection process. As a  
25 result, methods that have proven successful for *Agrobacterium*-mediated transformation of some monocots, such as rice and maize, have not been successful for transforming turfgrass.

An object of the present invention is to  
30 develop an efficient and reliable transformation system for turfgrass, mediated by *Agrobacterium tumefaciens*. Another object of the invention is to regenerate transgenic plants containing one or more foreign genes introduced by *Agrobacterium tumefaciens*-mediated  
35 transformation.

- 3 -

**SUMMARY OF THE INVENTION**

In accordance with the objects of the present invention, an *Agrobacterium*-mediated transformation system for turfgrass is provided, which is efficient and reliable. The invention also provides for the development of transgenic plants containing one or more transgenes of significant practical utility.

According to one aspect of the invention, a method of producing a transgenic turfgrass plant is provided. The method comprises: (a) providing regenerable callus tissue from the turfgrass plant; (b) inoculating the tissue with *Agrobacterium* carrying at least one vector for transformation, the vector comprising virulence genes that confer strong infectivity to *Agrobacterium*, in which vector is inserted a heterologous DNA construct operably linked to a promoter from a monocotyledonous species, and a selectable marker gene conferring antibiotic resistance to transformed cells operably linked to a promoter from a monocotyledonous species; (c) culturing the inoculated tissue under conditions that enable the *Agrobacterium* vector to transform cells of the tissue; (d) selectively culturing the inoculated tissue on a selection medium comprising the antibiotic; and (e) regenerating a transformed turfgrass plant from the selectively cultured tissue.

Preferably, the turfgrass is a species selected from the group consisting of creeping bentgrass, tall fescue, velvet bentgrass, perennial ryegrass, hard fescue, Chewings fescue, strong creeping fescue, colonial bentgrass and Kentucky bluegrass. In another preferred embodiment, the *Agrobacterium* comprises a binary vector system and the virulence genes therein are obtained from a plasmid within *Agrobacterium tumefaciens* strain 281. The promoter is preferably selected from the group

- 4 -

consisting of maize ubiquitin gene promoters, rice actin gene promoters, maize *Adh 1* gene promoters, rice or maize tubulin (Tub A, B or C) gene promoters, and alfalfa *His 3* gene promoters. The selectable marker gene preferably  
5 confers hygromycin resistance on transformed tissue. The callus used for the transformation preferably is obtained by culturing seeds of the turfgrass on a de-differentiation medium.

Also provided in accordance with another aspect  
10 of the invention is a transgenic turfgrass plant prepared by the aforementioned method. Seeds of the transgenic plant are provided as well. In preferred embodiments, the transgenic turfgrass plant comprises a transgene selected from the group consisting of genes encoding  
15 glucose oxidase, citrate synthase,  $\Delta$ -9 desaturase from *Saccharomyces cerevisiae* or *Cryptococcus curvatus*,  $\Delta$ -11 desaturase, a plant homolog of the neutrophil NADPH oxidase, a bacteriopsin from *Halobacterium halobium*, or pokeweed antiviral protein.

According to another aspect of the invention, a  
20 superbinary vector for *Agrobacterium*-mediated transformation of turfgrass is provided. The vector comprises: (a) a virulence region from a Ti plasmid of an *A. tumefaciens* strain that confers to the strain as  
25 strong a virulence as that displayed by *A. tumefaciens* strain 281; (b) a selectable marker gene operably linked to a promoter obtained from a gene of a monocotyledenous plant; and (c) a site for insertion of at least one  
30 additional coding sequence, operably linked to a promoter obtained from a gene of a monocotyledenous plant, the promoter being the same as or different from the promoter operably linked to the selectable marker gene. In preferred embodiments the virulence region is obtained from *Agrobacterium* strain 281. The promoter is selected

- 5 -

from the group consisting of maize ubiquitin gene promoters, rice actin gene promoters, maize *Adh 1* gene promoters, rice or maize tubulin (*Tub A*, *B* or *C*) gene promoters, and alfalfa *His 3* gene promoters. The  
5 selectable marker gene preferably confers hygromycin resistance on transformed cells. The site for insertion of the additional coding sequence preferably comprises a coding sequence of a reporter gene, and/or comprises a coding sequence of a gene encoding the useful proteins  
10 listed above.

According to another aspect of the invention, a turfgrass plant cell transformed with the aforementioned *Agrobacterium* vector is provided. Preferably the  
turfgrass is creeping bentgrass, tall fescue, velvet  
15 bentgrass, perennial ryegrass, hard fescue, Chewings fescue, strong creeping fescue, colonial bentgrass or Kentucky bluegrass. A transgenic turfgrass plant regenerated from the aforementioned transformed cell is also provided, as are seeds of the transgenic turfgrass  
20 plant.

Other features and advantages of the present invention will be understood by reference to the detailed description and examples that follow.

25

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions

Various terms used throughout the specification and claims to describe the invention. Unless otherwise  
30 specified, these terms are defined as set forth below.

With reference to nucleic acid molecules, the term "isolated nucleic acid" is sometimes used. This term, when applied to DNA, refers to a DNA molecule that is separated from sequences with which it is immediately

- 6 -

contiguous (in the 5' and 3' directions) in the naturally occurring genome of the organism from which it was derived. For example, the "isolated nucleic acid" may comprise a DNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the genomic DNA of a prokaryote or eukaryote. An "isolated nucleic acid molecule" may also comprise a cDNA molecule.

With respect to RNA molecules, the term "isolated nucleic acid" primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

The term "substantially pure" refers to a preparation comprising at least 50-60% by weight the compound of interest (e.g., nucleic acid, oligonucleotide, protein, etc.). More preferably, the preparation comprises at least 75% by weight, and most preferably 90-99% by weight, the compound of interest. Purity is measured by methods appropriate for the compound of interest (e.g. chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

When used herein in describing components of media or other experimental results, the term "about" means within a margin of commonly acceptable error for the determination being made, using standard methods. For tissue culture media in particular, persons skilled in the art would appreciate that the concentrations of various components initially added to culture media may

- 7 -

change somewhat during use of the media, e.g., by evaporation of liquid from the medium or by condensation onto the medium. Moreover, it is understood that the precise concentrations of the macronutrients, vitamins  
5 and carbon sources are less critical to the efficacy of the media than are the micronutrient, hormone and antibiotic concentrations.

Nucleic acid sequences and amino acid sequences can be compared using computer programs that align the  
10 similar sequences of the nucleic or amino acids thus define the differences. In the comparisons made in the present invention, the CLUSTLW program and parameters employed therein were utilized (Thompson et al., 1994, *supra*). However, equivalent alignments and  
15 similarity/identity assessments can be obtained through the use of any standard alignment software. For instance, the BLAST programs used to query sequence similarity in GenBank and other public databases may be used. The GCG Wisconsin Package version 9.1, available  
20 from the Genetics Computer Group in Madison, Wisconsin, and the default parameters used (gap creation penalty=12, gap extension penalty=4) by that program may also be used to compare sequence identity and similarity.

The term "substantially the same" refers to  
25 nucleic acid or amino acid sequences having sequence variations that do not materially affect the functionality of *cis* acting regulatory sequences (e.g, promoters, transcriptional response elements, etc.) or the nature of the encoded gene product (i.e. the  
30 structure, stability characteristics, substrate specificity and/or biological activity of the protein). With particular reference to nucleic acid sequences, the term "substantially the same" is intended to refer to

- 8 -

conserved sequences governing expression and to the coding region (referring primarily to degenerate codons encoding the same amino acid, or alternate codons encoding conservative substitute amino acids in the encoded polypeptide).

The terms "percent identical" and "percent similar" are also used herein in comparisons among nucleic acid sequences. When referring to nucleic acid molecules, "percent identical" refers to the percent of the nucleotides of the subject nucleic acid sequence that have been matched to identical nucleotides by a sequence analysis program. When referring to amino acid sequences, "percent identical" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical amino acids in the compared amino acid sequence by a sequence analysis program. "Percent similar" refers to the percent of the amino acids of the subject amino acid sequence that have been matched to identical or conserved amino acids. Conserved amino acids are those which differ in structure but are similar in physical properties such that the exchange of one for another would not appreciably change the tertiary structure of the resulting protein. Conservative substitutions are defined in Taylor (1986, J. Theor. Biol. 119:205).

With respect to oligonucleotides or other single-stranded nucleic acid molecules, the term "specifically hybridizing" refers to the association between two single-stranded nucleic acid molecules of sufficiently complementary sequence to permit such hybridization under pre-determined conditions generally used in the art (sometimes termed "substantially complementary"). In particular, the term refers to

- 9 -

hybridization of an oligonucleotide with a substantially complementary sequence contained within a single-stranded DNA or RNA molecule, to the substantial exclusion of hybridization of the oligonucleotide with single-stranded nucleic acids of non-complementary sequence.

A "coding sequence" or "coding region" refers to a nucleic acid molecule having sequence information necessary to produce a gene product, when the sequence is expressed.

The term "operably linked" or "operably inserted" means that the regulatory sequences necessary for expression of the coding sequence are placed in a nucleic acid molecule in the appropriate positions relative to the coding sequence so as to enable expression of the coding sequence. This same definition is sometimes applied to the arrangement other transcription control elements (e.g. enhancers) in an expression vector.

When describing the organization of a nucleic acid molecule, the term "upstream" refers to the 5' direction and the term "downstream" refers to the 3' direction.

The term "reporter gene" refers to a nucleic acid coding sequence that encodes a readily detectable gene product, which may be operably linked to a promoter region to form a chimeric gene, such that expression of the coding sequence is regulated by the promoter and the product of the coding sequence is readily assayed.

The term "selectable marker gene" refers to a gene that when expressed confers a selectable phenotype, such as antibiotic resistance, on a transformed cell or plant.

Transcriptional and translational control sequences, sometimes referred to herein as "expression

- 10 -

control" sequences or elements, or "expression  
regulating" sequences or elements, are DNA regulatory  
elements such as promoters, enhancers, ribosome binding  
sites, polyadenylation signals, terminators, and the  
5 like, that provide for the expression of a coding  
sequence in a host cell. The term "expression" is  
intended to include transcription of DNA and translation  
of the mRNA transcript.

The terms "promoter", "promoter region" or  
10 "promoter sequence" refer generally to transcriptional  
regulatory regions of a gene, which may be found at the  
5' or 3' side of the coding region, or within the coding  
region, or within introns. Typically, a promoter is a  
DNA regulatory region capable of binding RNA polymerase  
15 in a cell and initiating transcription of a downstream  
(3' direction) coding sequence. The typical 5' promoter  
sequence is bounded at its 3' terminus by the  
transcription initiation site and extends upstream (5'  
direction) to include the minimum number of bases or  
20 elements necessary to initiate transcription at levels  
detectable above background. Within the promoter  
sequence is a transcription initiation site (conveniently  
defined by mapping with nuclease S1), as well as protein  
binding domains (consensus sequences) responsible for the  
25 binding of RNA polymerase.

A "vector" is a replicon, such as plasmid,  
phage, cosmid, or virus to which another nucleic acid  
segment may be operably inserted so as to bring about the  
replication or expression of the segment.

30 The term "nucleic acid construct" or "DNA  
construct" refers to genetic sequence used to transform  
plant cells and generate progeny transgenic plants. At  
minimum a DNA construct comprises a coding region for a

- 11 -

selected gene product, operably linked to 5' and 3' regulatory sequences for expression in transformed plants. In preferred embodiments, such constructs are chimeric, i.e., the coding sequence is from a different source one or more of the regulatory sequences (e.g., coding sequence from tobacco and promoter from maize). However, non-chimeric DNA constructs also can be used. In addition to methods specifically described herein, the transforming DNA may be prepared according to standard protocols such as those set forth in Ausubel et al. (1999). A plant species or cultivar may be transformed with a DNA construct (chimeric or non-chimeric) that encodes a polypeptide from a different plant species or cultivar, or a non-plant species. Alternatively, a plant species or cultivar may be transformed with a DNA construct (chimeric or non-chimeric) that encodes a polypeptide from the same plant species or cultivar. The term "transgene" is sometimes used to refer to the DNA construct within the transformed cell or plant.

20 A cell has been "transformed" or "transfected" by a DNA construct when such DNA has been introduced inside the cell. The transforming DNA may or may not be integrated (covalently linked) into the genome of the cell. For example, the transforming DNA may be maintained on an episomal element such as a plasmid. With respect to eukaryotic cells, a stably transformed cell is one in which the transforming DNA has become integrated into a chromosome so that it is inherited by daughter cells through chromosome replication. This stability is demonstrated by the ability of the eukaryotic cell to establish cell lines or clones comprised of a population of daughter cells containing the transforming DNA. A "clone" is a population of cells derived from a single cell or common ancestor by mitosis.

- 12 -

A "cell line" is a clone of a primary cell that is capable of stable growth *in vitro* for many generations.

## II. Description

5           The present invention provides an efficient and reliable transformation system for turfgrass species. The species chosen to develop this system uses creeping bentgrass, *Agrostis palustris* Huds. Due to the reasonably close similarity among various turfgrasses  
10 (i.e., in physiology, genome organization, etc.), this transformation system will have broad applicability to many different kinds of turfgrass. For instance, the system also has been applied to velvet bentgrass, *Agrostis canina* L. and tall fescue, *Festuca arundinacea*  
15 Scheb., and although the transformation efficiency is not as great as for creeping bentgrass, it is clear that transformants can be obtained from several species. Other turfgrasses contemplated for the transformation system of the invention include, but are not limited to,  
20 perennial ryegrass, hard fescue, Chewings fescue, strong creeping fescue, colonial bentgrass and Kentucky bluegrass.

Binary vectors are commonly used in *Agrobacterium*-mediated transformation. Recently, a  
25 "superbinary" vector system was developed (Komari et al., 1990; Saito et al., 1992; Hiei et al., 1994; U.S. Patent No. 5,731,139 to Komari et al.). In this system the plasmid that carries the T-DNA also contains certain virulence genes from strain A281, which is known for high  
30 efficiency of transformation.

In arriving at the present invention, the conditions and vectors described for transformation of rice (Hiei et al., 1994) and maize (Ishida et al., 1996) were initially used, but without success in creeping  
35 bentgrass. It was suspected that a major problem with

- 13 -

these vectors was the CaMV35S promoter which drives the expression of the selectable marker in the rice and maize systems. Therefore, a new vector was constructed, which has the components of the "superbinary vector" developed by Hiei et al. (1994), coupled with a suitable selectable marker (hygromycin resistance) and the easily scorable  $\beta$ -glucuronidase (GUS) reporter gene. More significantly, in the new vector, the expression of each of these respective transgenes is driven by promoters for strong, constitutive expression in monocots, e.g., the rice actin and maize ubiquitin promoters. Independent transformations, conducted over the course of several months, demonstrated that an *Agrobacterium tumefaciens* such as this can indeed transfer DNA into the chromosomes of various turfgrasses.

Certain features of the *Agrobacterium*-mediated turfgrass transformation system describe herein are believed to be particularly important for successful transformation and regeneration of transgenic turfgrass. One of these is the use of Ti plasmids from strongly infective *Agrobacterium* strains. Preferred for use are superbinary hybrid *Agrobacterium* vectors, such as pSB1 and pSB11 (Komari et al., 1996). These and other superbinary vectors that can be modified for use in the present invention are described in U.S. Patent No. 5,731,179 to Komari et al.. These vectors contain a DNA region containing *virB* gene and *virG* gene in virulence region of Ti plasmid pTiBo542 of *Agrobacterium tumefaciens*, which is a Ti plasmid contained in *A. tumefaciens* strain A281 (ATCC Accession No. 37349), a strain known for its strong virulence. The *virB* and *virG* genes in the virulence region of pTiBo542 are contained a 15.2 kb KpnI restriction fragment, which itself may be used in the present invention. Moreover, although vectors comprising the hypervirulence-conferring portions

- 14 -

of pTiBo542 are exemplified herein, it will be appreciated by persons skilled in the art that corresponding genes from any highly virulent *Agrobacterium* strain may be used instead. These include strains that are presently available as well as strains that may be discovered in the future.

A particularly important feature of the invention is the modification of *Agrobacterium* vectors to contain promoters and other regulatory sequences particularly suited for expression in turfgrasses. Thus, the strong constitutive promoters from the rice actin gene and the maize ubiquitin gene are exemplified herein. However, any constitutively expressed or inducible promoter for expression in monocots may be used in the present invention. Examples of other constitutive promoters suitable for use in the present invention are the maize *Adh 1* promoter, the rice or maize tubulin (*Tub A*, *B* or *C*) promoters, and the alfalfa *His 3* promoter. Monocotyledenous tissue-specific promoters also may be utilized. For example, seed-specific promoters suitable for use in the present invention include zein promoters, such as the 27-kDa zein promoter or the 10-kDa zein promoter.

The choice of selectable marker is also important. Hygromycin is exemplified herein as a selectable marker. However, other selectable markers are also suitable for use in the invention. For instance, resistance to phosphothricin herbicides is particularly useful for selecting transformed monocotyledenous plant cells. In addition, kanamycin is routinely used as a selection medium for plant transformation, though it is less preferred for use in the present invention than is hygromycin.

For *Agrobacterium*-mediated transformation, an antibiotic to eliminate *Agrobacterium* is included in the

- 15 -

selection medium. Cefotaxime is the preferred antibiotic for this purpose. However, Augmentin (amoxicillin and lithium clavulenate) and carbenicillin also have been found effective for eliminating *Agrobacterium*.

5           If a reporter gene is utilized, it can be any one of several commonly used in the art. Examples of suitable reporter genes include, but are not limited to, genes encoding  $\beta$ -glucuronidase (GUS), luciferase, chloramphenicol acetyl transferase (CAT), green  
10   fluorescent protein (GFP) and modified forms of GFP (e.g., EGFP, EPFP and GFP<sub>uv</sub>).

          In addition, the use of friable, regenerable callus for the transformation protocol is considered particularly important to successful, efficient  
15   transformation and regeneration of intact plants. In a preferred embodiment, the callus is generated from embryogenic tissue, most preferably mature seeds, though it may also be generated from seed parts or immature seeds. Other organogenic tissues may also be utilized as  
20   starting material for callus growth. It has been found in accordance with the present invention that the culture media described herein will facilitate the growth of appropriate (i.e. friable, regenerable) callus tissue, which is best harvested as soon as a sufficient amount of  
25   callus is generated from the starting material.

          To optimize the conditions for efficient transformation of turfgrass, a number of parameters may be systematically altered to determine the best conditions to achieve efficient and easily reproducible  
30   transformation. Conditions that may be altered include: 1) the culture medium and induction agents immediately prior to and during co-cultivation with *Agrobacterium*; 2) inoculation and co-culture methods and time period; 3) the presence and absence of surfactants in the

- 16 -

inoculation medium; and 4) use of embryogenic callus versus suspension cell cultures. Optimum conditions have been established for creeping bentgrass. Modifications of these conditions have enabled the transformation of callus obtained from velvet bentgrass and tall fescue.

Thus, the present invention provides a method for generating transgenic turfgrass using *Agrobacterium* vectors, that heretofore has been unavailable. A preferred embodiment of the present invention comprises the following transformation/regeneration protocol, based on *Agrobacterium*-mediated transformation.

1) The starting tissue (e.g., mature seeds in a particularly preferred embodiment) of the selected turfgrass are surface-sterilized and placed on a standard de-differentiation tissue culture medium (see the examples) for about 3-6 weeks, preferably in the dark, at room temperature.

2) Proliferating calli are selected and transferred to fresh medium of the same type on a regular basis. Only callus that is friable and regenerable should be selected for further culture.

3) Prior to exposure to *Agrobacterium*, the chosen callus is transferred to new medium to promote active cell division; the callus is used for transformation within about a week of the final transfer to fresh medium.

4) The *Agrobacterium* strain carrying the transforming plasmids is grown, induced with acetosyringone (a phenolic compound demonstrated to induce *Agrobacterium vir* gene expression) and resuspended in an inoculation medium comprising acetosyringone and, optionally, a surfactant such as pluronic F-68 or another suitable surfactant.

5) Optionally, callus may be pre-treated prior to inoculation with *Agrobacterium* by vacuum-infiltration

- 17 -

with inoculation medium containing acetosyringone.

6) The callus tissue is then placed in the presence of the *Agrobacterium* suspension to allow the bacteria to infiltrate the callus tissue. Optionally, excess *Agrobacterium* may thereafter be removed from the calli by gentle vacuum filtration. Calli, on a sterile filter, are then placed on co-cultivation solid medium (standard de-differentiation medium but containing glucose and acetosyringone). Co-cultivation of the calli with the *Agrobacterium* cells that remain associated with the calli is allowed to proceed for a few (e.g., three) days, the object being to allow sufficient time for penetration of the T-DNA into the plant cells while avoiding overgrowth of the calli with the *Agrobacterium*.

7) The calli are then transferred to selection medium containing the selection antibiotic (e.g., hygromycin) and the antibiotic for removal of the *Agrobacterium*. Calli are kept on this medium for several weeks (e.g. 6-8 weeks) and checked periodically for proliferation of the calli on the selection medium.

8) New growth that appears on calli on the selection medium is first transferred to fresh selection medium and allowed to proliferate. This first selection is best performed when the new growth is as small as possible in order to ensure that independent transformants are selected and proliferated in the absence of other independent transformants.

9) After sufficient proliferation, a small portion of each of the putatively transformed calli is tested for expression of the gene of interest, and/or for expression of a detectable marker gene (e.g., GUS activity). The remaining portions of positive-testing calli are retained on the selection medium for continued subculture and proliferation. Subsequently, the transformed calli are transferred to a regeneration

- 18 -

medium containing growth regulators to promote shoot differentiation.

10) When sufficiently developed, the shoots are transferred to a second regeneration medium formulated to further stimulate root growth. After a sufficient growth phase, plantlets are transferred to either new tissue culture medium or to soil or equivalent planting media.

This invention provides transgenic turfgrass produced by the above-described methods, and also is intended to encompass cells and tissues of those plants, including, but not limited to, leaves, stems, shoots, roots, flowers, fruits and seeds. In a preferred embodiment, seeds of the transgenic plants produced by the methods of the invention are provided.

The plants grown from the aforementioned seeds, or seeds from other turfgrass species or varieties, or the progeny thereof, all of which are considered within the scope of this invention, are used in crosses and selection methods to transfer genes of interest into other turfgrass genotypes, cultivars, varieties and the like.

Plants grown from transgenic turfgrass seeds can also be used to detect the presence of the inserted transgene and vector sequences using DNA extraction, cleavage by one or more restriction endonucleases, and analysis, e.g., Southern blotting using probes derived from the gene or genes of interest. In this manner, the transfer of foreign genes into progeny of breeding crosses can be monitored.

The *Agrobacterium*-mediated turfgrass transformation system of the present invention can be used to introduce many genes of interest into different turfgrass species or varieties. Accordingly, the present invention provides several specific hybrid vectors for *Agrobacterium*-mediated transformation of turfgrass. A

- 19 -

number of gene constructs are of considerable practical utility as used to create one or more different transgenic turfgrasses. These include: 1) the gene encoding glucose oxidase from *Aspergillus niger* which when expressed in potato provides resistance to bacterial and fungal pathogens through its production of  $H_2O_2$  in the plant apoplast (Wu et al., 1995); 2) the gene encoding citrate synthase from *Pseudomonas aeruginosa* which when expressed in the cytoplasm of tobacco provides tolerance to toxic levels of aluminum in the soil (Manuel et al., 1997); 3) the genes encoding  $\Delta$ -9 desaturase from *Saccharomyces cerevisiae* (Stukey et al., 1990) and *Cryptococcus curvatus* (Meesters et al., 1997) and  $\Delta$ -11 desaturase from *Trichoplosia ni* (Knipple et al., 1998). When the *ole1* gene from yeast was expressed in tomato and eggplant, it provided resistance to fungal pathogens (Wang et al., 1998); 4) the recently identified gene encoding a plant homolog of the neutrophil NADPH oxidase which is thought to be responsible for the oxidative burst which is critical in plant defense against pathogens (Keller et al., 1998); 5) the gene encoding bacteriopsin, a proton pump from the bacterium, *Halobacterium halobium*, which has been demonstrated to protect transgenic tobacco expressing the protein against plant pathogens (Mittler et al., 1995); and 6) the gene encoding pokeweed antiviral protein, which is a ribosome-inactivating protein from the plant *Phytolacca americana*; expression of this gene in tobacco provides resistance to viral and fungal pathogens (Lodge et al., 1993; Zoubenko et al., 1997).

The following examples are provided to describe the invention in greater detail. They are intended to illustrate, not to limit, the invention.

- 20 -

## EXAMPLE 1

**Construction of *Agrobacterium tumefaciens* Strain LBA4404, containing pSB111SH Superbinary Hybrid Vector**

5           The vector pTOK233 described by Hiei et al. (1994) for transformation of rice was not useful for transformation of creeping bentgrass despite numerous attempts. It was suspected that the cauliflower mosaic 35S promoter was the source of the problem in that it appears to be a very weak promoter in turfgrass. Accordingly, an alternate pair of plasmid vectors was obtained, namely pSB1 and pSB11 (Japan Tobacco, Inc. Plant Breeding and Genetics Research Laboratory, 700 Higashibara, Toyoda, Iwata, Shizuoka, Japan). pSB11 is a 15 6.3 kb plasmid which contains a multicloning region with a number of unique restriction sites between the left and right border sequences delineating the T-DNA region that will be transferred from *Agrobacterium tumefaciens* to the plant genome. Monocot "expression cassettes" expressing 20 hygromycin B phosphotransferase (to confer hygromycin resistance) and  $\beta$ -glucuronidase (GUS, as a scorable reporter) were inserted into pSB11 into the multiple cloning region between the T-DNA border sequences. The following strategy was used.

25           First, plasmid pAch1 (provided by Dr. German Spangenberg) was partially digested with *Sac* I which yielded several restriction fragments, of which the 2645 bp fragment was gel-purified. This fragment contains a 3' modified *hph* coding sequence (conferring hygromycin 30 resistance) cloned inframe and downstream from the Act 1 5' promoter sequence, first exon (non-coding), first intron and a portion of the second, ATG-containing exon of the gene. Transcription termination signals are provided by the CaMV35S terminator. This 2645 bp 35 fragment was cloned into the unique *Sac* I restriction site between the right and left border sequences of

- 21 -

plasmid pSB11. This plasmid was named pSB11S. Next, a 4175 bp *Hind* III fragment was excised from plasmid pAHC25, kindly provided by Dr. Peter Quail. This fragment contains the promoter, 5' untranslated exon, and first intron of the maize ubiquitin (*Ubi1*) gene fused with the GUS reporter gene coding sequence (derived from pBI101.2). This *Hind* III restriction fragment was cloned into the unique *Hind* III site of pSB11S, between the right and left border sequences of the plasmid. The resultant plasmid was named pSB11SH.

Next, this intermediate vector (pSB11SH) was electroporated into *Agrobacterium tumefaciens* LBA4404 containing the "acceptor vector" pSB1. Plasmid pSB1 is a "superbinary vector" which carries a 15.2 kb *Kpn* I fragment containing additional copies of the *vir* B, *vir* C and *vir* G virulence genes to enable efficient T-DNA transfer to monocots. Following homologous recombination between the "acceptor vector" and the "intermediate vector", a co-integrate "hybrid vector" was obtained. Bacteria expressing the drug resistance markers derived from both the acceptor and intermediate vectors were selected on AB plates containing 10  $\mu$ g/ml tetracycline and 50  $\mu$ g/ml spectinomycin, and the resultant strains carrying the "hybrid vector" were used in plant transformations.

## EXAMPLE 2

### Agrobacterium-Mediated Transformation of Creeping Bentgrass With Superbinary Hybrid Vector pSB11SH

In this example, an *Agrobacterium* transformation system for creeping bentgrass, *Agrostis palustris*, is described, which utilizes the superbinary hybrid test vector described in Example 1.

- 22 -

**Media:****MMSG Medium****Per Liter:**

5	Murashige and Skoog basal salts	4.33 g
	sucrose	30 g
	Gamborg vitamins (1000x)	1 ml
	casein hydrolysate	500 mg
	dicamba (dichloro-o-anisic acid)	6.6 mg
10	6 BAP (6-benzylaminopurine)	0.5 mg
	Adjust the pH to 5.6 to 5.8.	
	Add GEL-GRO™*	2.4 g
	Autoclave and dispense into Petri dishes.	
	(*GEL-GRO™ is gellan gum, an agar substitute,	
15	manufactured by ICN Biomedicals, Inc. If GEL-GRO™ is	
	unavailable, agar or an equivalent agar substitute may be	
	utilized.)	

**Modified AAM Medium****Per Liter:**

20	glutamine	875 mg
	aspartic acid	266 mg
	arginine	174 mg
	glycine	7.5 mg
25	sucrose	88.5 mg
	2,4-D	1 mg
	kinetin	0.2 mg
	GA <sub>3</sub>	0.1 mg
	Murashige & Skoog vitamins (1000x)	1 ml
30	casamino acids	500 mg
	glucose	36 g
	Adjust pH to 5.2 and autoclave.	
	Add acetosyringone to a final concentration of 100 µM.	

- 23 -

**Regeneration Medium MSO I****Per Liter:**

	Murashige & Skoog basal salts	4.33 g
	Gamborg vitamins (1000x)	1 ml
5	sucrose	30 g
	myo-inositol	100 mg
	6-benzylaminopurine	1 mg
	Adjust pH to 5.8.	
	Add GEL-GRO™	2.4 g
10	Autoclave and dispense into plates.	

**Regeneration Medium MSO II****Per Liter:**

	Murashige & Skoog basal salts	4.33 g
15	Gamborg vitamins (1000x)	1 ml
	sucrose	30 g
	myo-inositol	100 mg
	Adjust pH to 5.8.	
	Add GEL-GRO™	2.4 g
20	Autoclave and dispense into petri dishes and plantcons.	

**AB medium**

1. Prepare a stock of AB Buffer at 20X and autoclave:

25 **Per Liter:**

$K_2HPO_4$	60 g
$NaH_2PO_4$	20 g

2. Prepare a stock of AB Salts at 20X and autoclave:

30 **Per Liter:**

	$NH_4Cl$	20 g
	$MgSO_4 \cdot 7H_2O$	6 g
	KCl	3 g
	$CaCl_2$	0.2 g
35	$FeSO_4 \cdot 7H_2O$	50 mg

- 24 -

3. For 1 liter of medium for plates:  
Mix 5 g of glucose, 15 g of agar, and 900 ml of H<sub>2</sub>O and autoclave.  
Add 50 ml of the 20X stock of AB Buffer and 50 ml of the 20X stock of AB Salts.  
Add 10 mg tetracycline and 50 mg spectinomycin.  
Pour into petri dishes.

**Production of regenerable callus:** Mature seeds of creeping bentgrass (Crenshaw cultivar) were surface-sterilized and plated on MMSG medium. The plates were kept in the dark at room temperature for 3-6 weeks. The proliferating calli were selected and transferred to new MMSG medium on a regular basis. Only callus that was friable and regenerable was used for transformation. The chosen callus was transferred to new MMSG medium prior to co-cultivation to promote active cell division and was used for transformation within a week after transferring to new plates. The nature of the callus (i.e., friability, regenerability and active growth) is believed to play a key role in obtaining efficient transformation.

**Induction of *Agrobacterium tumefaciens* with acetosyringone:** *Agrobacterium tumefaciens* LBA4404, harboring vector pSB111SH, was streaked from a glycerol stock stored at -80°C and grown at 28°C on plates containing AB medium, supplemented with 10 µg/ml tetracycline and 50 µg/ml spectinomycin. After three to six days, the cells were scraped from the plate and suspended in modified AAM medium containing 100 µM acetosyringone to an OD<sub>660</sub> of approximately 0.5. The bacterial suspension was left at 25°C in the dark with shaking for 3.5 hours before using it for co-cultivation.

**Co-cultivation:** A given amount of friable callus was mixed with the pre-induced *Agrobacterium*

- 25 -

suspension and incubated at room temperature in the dark for 1.5 hours. Then the contents were poured into a sterile Buchner-funnel containing a sterile Whatman filter paper. Mild vacuum was applied to drain the excess *Agrobacterium* suspension. Then, the filter was moved to a plate containing MMSG medium supplemented with 100  $\mu$ M acetosyringone, and the plate was stored in the dark at room temperature for three days.

**Selection and regeneration of transformants:**

Subsequent to the three day co-cultivation, the co-cultivated calli were rinsed with 250  $\mu$ g/ml cefotaxime to suppress bacterial growth, and the calli were placed on agar plates containing MMSG medium containing 200  $\mu$ g/ml hygromycin and 250  $\mu$ g/ml cefotaxime. The calli were kept in the dark at room temperature for 6-8 weeks and checked periodically for proliferation of the calli on hygromycin.

Subsequently, the hygromycin-resistant calli were placed on regeneration medium containing hygromycin and cefotaxime. The proliferating calli were first moved to Regeneration Medium I (MSO I) containing cefotaxime and hygromycin. These calli were kept in the dark at room temperature for a week and were then moved to light for approximately two weeks. The tiny plants were separated and transferred to deep petri plates containing Regeneration Medium II (MSO II) to promote root growth. Hygromycin and cefotaxime were included in the medium to respectively maintain selection pressure and kill any remaining *Agrobacterium* cells. After 2-3 weeks, or when the plants were 1.5-2 cm tall, they were moved to plantcons containing MSO II without antibiotics. When the plants were ~10 cm tall and had developed extensive root systems, they were transferred to soil and grown in the laboratory for 3-4 weeks with 12 hours light/day. The plants were transferred to 6" pots to the greenhouse,

- 26 -

where the temperature was maintained between 21° - 25°C. Supplemental lighting added approximately 50  $\mu\text{E m}^{-2} \text{ sec}^{-1}$  at canopy level when natural light was low and provided a minimal light period of 14 hours.

5 In connection with the transformation of creeping bentgrass, it should be noted that the media and protocols described below in Examples 3 and 4 may also be used to successfully transform and regenerate creeping bentgrass.

10

### EXAMPLE 3

#### Agrobacterium-Mediated Transformation of Tall Fescue With Superbinary Hybrid Vector pSB111SH

15 In this example, an *Agrobacterium* transformation system for tall fescue, *Festuca arundinacea* Schreb., is described, which utilizes the superbinary hybrid test vector described in Example 1.

#### Media:

20

#### MMSG Medium

This medium was formulated as described in Example 2.

#### Inoculation Medium

#### Per Liter:

25	Murashige & Skoog basal salts	440 mg
	sucrose	30 g
	dicamba	6.6 mg
	6 BAP (6-benzyl aminopurine)	0.5mg
	Gamborg vitamins (1000x)	1ml
30	casein hydrolysate	500 mg
	glucose	10 g

Adjust pH to 5.7 and autoclave.

Immediately before use add acetosyringone to a final concentration of 200  $\mu\text{M}$  and pluronic F-68 to a final

35 concentration of 0.02%.

- 27 -

**Co-Cultivation Medium****Per Liter:**

	Murashige and Skoog basal salts	4.4 g
	sucrose	30 g
5	glucose	10 g
	Gamborg vitamins (1000x)	1 ml
	casein hydrolysate	500 mg
	dicamba (dichloro-o-anisic acid)	6.6 mg
	6 BAP (6-benzylaminopurine)	0.5 mg
10	Adjust the pH to 5.7	
	Add GEL-GRO™	2.4 g
	Autoclave	
	Add acetosyringone	39.2 mg

15 **Regeneration Medium MSO I**

This medium was formulated and prepared as described in Example 2.

**Regeneration Medium MSO II**

20 This medium was formulated and prepared as described in Example 2.

**AB medium**

This medium was formulated and prepared as described in Example 2.

25 **Production of regenerable callus:** Mature seeds of tall fescue were surface-sterilized and plated on MMSG medium. The plates were kept in the dark at room temperature for 3-6 weeks. The proliferating calli were selected and transferred to new MMSG medium on a regular basis. Callus chosen for transformation was transferred to new MMSG medium prior to co-cultivation to promote active cell division. The nature of the callus (i.e., friability, regenerability and active growth) is believed to play a key role in obtaining efficient transformation.

30

- 28 -

**Preparation of *Agrobacterium tumefaciens***

**suspension:** *Agrobacterium tumefaciens* LBA4404, harboring vector pSB111SH, was streaked from a glycerol stock stored at -80°C and grown at 28°C on plates containing AB medium, supplemented with 10 µg/ml tetracycline and 50 µg/ml spectinomycin. After three to six days, the cells were scraped from the plate and suspended in inoculation medium containing 200 µM acetosyringone and 0.02% pluronic F-68 to an OD<sub>660</sub> between 0.5 and 0.8.

**10 Co-cultivation of callus with *Agrobacterium*:**

The friable callus chosen for transformation was placed in a sterile tube and mixed with 30 ml of *Agrobacterium* suspension. The tube was capped and covered with aluminum foil, and the contents were mixed by inversion and gently shaken for about 1.5 hr. The contents of the tube were poured into a Buchner funnel, fitted with Whatman filter. Mild vacuum was applied to flask. The filter disks containing callus with *Agrobacterium* were moved to co-cultivation plates containing acetosyringone (200 µM) and glucose at 10 g/l. The plates were sealed with parafilm and placed in the dark at room temperature for three days.

**Selection and regeneration of transformants:**

Subsequent to the three day co-cultivation, the co-cultivated calli were rinsed with 250 µg/ml cefotaxime solution to suppress bacterial growth, and the calli were placed on MMSG medium containing 200 µg/ml hygromycin and 250 µg/ml cefotaxime. The calli were kept in the dark at room temperature for 6-8 weeks and checked periodically for proliferation of the calli on hygromycin. The hygromycin-resistant calli were moved to new MMSG plates with hygromycin and kept in the dark at room temperature until well proliferated. Then a portion of the hygromycin-resistant callus was tested for GUS activity

- 29 -

to ensure that transformation had occurred.

Portions of the putatively transformed calli were then moved to Regeneration Medium I (MSO I) containing cefotaxime and hygromycin and kept in the dark room at room temperature for a week and were subsequently moved to the light for regeneration. The tiny shoots were separated and transferred to deep petri plates containing Regeneration Medium II (MSO II) to promote root growth and hygromycin and cefotaxime to maintain respectively selection of the transformants and kill any remaining *Agrobacterium*.

#### EXAMPLE 4

##### **Agrobacterium-Mediated Transformation of Velvet Bentgrass With Superbinary Hybrid Vector pSB111SH**

In this example, an *Agrobacterium* transformation system for velvet bentgrass, *Agrostis canina* L., is described, which utilizes the superbinary hybrid test vector described in Example 1.

**Media:** All media were formulated and prepared as described in Example 3.

**Production of regenerable callus:** Mature seeds of velvet bentgrass were surface-sterilized and plated on MMSG medium. The plates were kept in the dark at room temperature for 3-6 weeks. The proliferating calli were selected and transferred to new MMSG medium on a regular basis. Callus chosen for transformation was transferred to new MMSG medium prior to co-cultivation to promote active cell division. The nature of the callus (*i.e.*, friability, regenerability and active growth) is believed to play a key role in obtaining efficient transformation.

- 30 -

**Preparation of *Agrobacterium tumefaciens***

**suspension:** *Agrobacterium tumefaciens* LBA4404, harboring vector pSB111SH, was streaked from a glycerol stock stored at -80°C and grown at 28°C on plates containing AB medium, supplemented with 10 µg/ml tetracycline and 50 µg/ml spectinomycin. After three to six days, the bacterial lawn was scraped from one (82 mm diameter) plate and suspended in 6 ml inoculation medium containing 200 µM acetosyringone. The bacterial suspension was left at 28°C in the dark with shaking overnight. In the morning acetosyringone was added to a final concentration of 400 µM and pluronic F-68 to 0.02%.

**Co-cultivation:** A sterile filter was placed on a co-cultivation plate containing acetosyringone (200 µM) and glucose (10 gm/l). Onto this filter was placed one large clump of friable callus which was then gently broken up and dispensed evenly over the filter. About 500 µl of *Agrobacterium* suspension was pipetted evenly onto the callus. The plates were sealed with parafilm and placed in the dark at room temperature for three days.

**Selection and regeneration of transformants:**

Subsequent to the three day co-cultivation, the co-cultivated calli were rinsed with 250 µg/ml cefotaxime solution to suppress bacterial growth, and the calli were placed on MMSG medium containing 200 µg/ml hygromycin and 250 µg/ml cefotaxime. The calli were kept in the dark at room temperature for 6-8 weeks and checked periodically for proliferation of the calli on hygromycin. The hygromycin-resistant calli were moved to new MMSG plates with hygromycin and kept in the dark at room temperature until well proliferated. Then a portion of the hygromycin-resistant callus was tested for GUS activity to ensure that transformation had occurred.

- 31 -

Portions of the putatively transformed calli were then moved to Regeneration Medium I (MSO I) containing cefotaxime and hygromycin and were kept in the dark room at room temperature. After one week, they were moved to the light for regeneration. The tiny shoots were separated and transferred to deep petri plates containing Regeneration Medium II (MSO II) to promote root growth and hygromycin and cefotaxime to maintain respectively selection pressure and kill any remaining *Agrobacterium* cells.

#### REFERENCES

- Aldemita, R.R. and T.K. Hodges (1996) *Agrobacterium tumefaciens*-mediated transformation of *japonica* and *indica* rice varieties. *Planta* 199:612-617.
- Belanger, F.C., C.L. Laramore, and P.R. Day (1996) Turfgrass biotechnology. *Rutgers Turfgrass Proceedings* 28:1-3.
- Cheng, M., J.E. Fry, S. Pang, H. Zhou, C.M. Hironaka, D.R. Duncan, T.W. Conner and Y. Wan (1997) Genetic transformation of wheat mediated by *Agrobacterium tumefaciens*. *Plant Physiology* 115:971-980.
- Czernilofsky, A.P., R. Hain, L. Herrera-Estrella, H. Lorz, E. Goyvaerls, B.J. Baker and J. Schell (1986) Fate of selectable marker DNA integrated into the genome of *Nicotiana tabacum*. *DNA* 5:101-113.
- Hiei, Y., S. Ohta, T. Komari and T. Kumashiro (1994) Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant Journal* 6:271-282.
- Ishida, Y., H. Saito, S. Ohta, Y. Hiei, T. Komari and T. Kumashiro (1996) High efficiency transformation of maize (*Zea mays* L.) mediated by *Agrobacterium tumefaciens*. *Nature Biotechnology* 14:745-

- 32 -

750.

Keller, T., H.G. Damude, D. Werner, P. Doerner, R.A. Dixon and C. Lamb (1998) A plant homolog of the neutrophil NADPH oxidase gp91<sup>phox</sup> subunit gene encodes a plasma membrane protein with Ca<sup>2+</sup> binding motifs. *Plant Cell* 10:255-266.

Knipple, D.C., C.L. Rosenfield, S.J. Miller, W. Liu, J. Tang, P.W.K. Ma and W.L. Roelofs (1998) Cloning and functional expression of a cDNA encoding a pheromone gland-specific acyl-CoA  $\Delta^{11}$ -desaturase of the cabbage looper moth, *Trichoplusia ni*. *Proc. Natl. Acad. Sci. USA* 95:15287-15292.

Komari, T. (1990) Transformation of cultured cells of *Chenopodium quinoa* by binary vectors that carry a fragment of DNA for the virulence region of pTiB0542. *Plant Cell Rep.* 9:303-306.

Komari, T., Y. Hiei, Y. Saito, N. Murai, and T. Kumashiro (1996) Vectors carrying two separate T-DNAs for co-transformation of higher plants mediated by *Agrobacterium tumefaciens* and segregation of transformants free from selection markers. *Plant J.* 10: 165-174.

Lee, L. (1996) Turfgrass biotechnology. *Plant Science* 115:1-8.

Lodge, J., W.K. Kaniewski and N.E. Tumer (1993) Broad spectrum virus resistance in transgenic plants expressing pokeweed antiviral proteins. *Proc. Natl. Acad. Sci. USA* 90: 7089-7093.

Manuel de la Fuente, J., V. Ramires-Rodriguez, J.L. Cabrera-Ponce and L. Herrera-Estrella (1997) Aluminum tolerance in transgenic plants by alteration of citrate synthesis. *Science* 276:1566-1568.

May, G.D., R. Afza, H.S. Mason, A. Wiecko, F.J. Novak and C.J. Arntun (1995) Generation of transgenic banana (*Musa acuminata*) plants via *Agrobacterium*-mediated transformation. *Biotechnology* 13:486-492.

- Meesters, P.A.E.P., J. Springer and G. Eggink (1997) Cloning and expression of the  $\Delta$ -9 fatty acid desaturase gene from *Cryptococcus curvatus* ATCC 20509 containing histidine boxes and a cytochrome  $b_5$  domain. *Appl. Microbiol. Biotechnol.* 47:663-667.
- 5 Mittler, R., V. Shulaev and E. Lam (1995) Coordinated activation of programmed cell death and defense mechanisms in transgenic tobacco plants expressing a bacterial proton pump. *Plant Cell* 7: 29-42.
- 10 Saito, Y. T. Komari, C. Matsuta, Y. Hayashi, T. Kumashiro and Y. Takanami (1992) Cucumber mosaic virus-tolerant transgenic tomato plants expressing satellite RNA. *Theor. App. Genet.* 83: 679-683.
- 15 Stucky, J.E., V.M. Mc Donough and C.E. Martin (1990) The *ole1* gene of *Saccharomyces cerevisiae* encodes the  $\Delta$ -9 fatty acid desaturase and can be functionally replaced by the rat stearyl-CoA desaturase gene. *J. Biol. Chem.* 265:20144-20149.
- 20 Tingay, S., D. McElroy, R. Kalla, S. Fieg, M. Wang, S. Thornton and R. Brettell (1997) *Agrobacterium tumefaciens*-mediated barley transformation. *Plant Journal* 11:1369-1376.
- 25 Wang, C., C.-K. Chin and A. Chen (1998) Expression of the yeast delta nine desaturase gene in tomato enhances its resistance to powdery mildew. *Physiological and Molecular Plant Pathology* 38:255-263.
- 30 Wu, G., B.J. Shortt, E.B. Lawrence, E.B. Levine, K.C. Fitzsimmons and D.M. Shah (1995) Disease resistance conferred by expression of a gene encoding  $H_2O_2$ -generating glucose oxidase in transgenic potato plants. *Plant Cell* 7:1357-1368.
- Yuan, R. (1998) Biotech driving innovations in rice cultivation. *Genetic Engineering News*, Jan. 15 issue: 13-32.

- 34 -

Zoubenko, O., F. Uckun, Y. Hur, I. Chet and N. Tumer (1997) Plant resistance to fungal infection induced by nontoxic pokeweed antiviral protein mutants. *Nature Biotech.* 15: 992-996.

5

While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be limited to such embodiments. Various  
10 modifications may be made thereto without departing from the scope and spirit of the present invention, as set forth in the following claims.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193